

## SHOUTS: 4 (see end)

---

no logo available

---

Wednesday, 23 May 2007

9:30 – 10:45 (1 course)

9:30 – 12:00 (2 courses)

---

EXAMINATION FOR THE DEGREES OF B.Sc. M.Sci. AND M.A. ON  
THE HONOURS STANDARD

---

Physics 3 – Chemical Physics 3 – Physics with Astrophysics 3  
Theoretical Physics 3M – Joint Physics 3

**P304D and P304H**

[ **PHYS3031 and PHYS4025** ]

# Quantum Mechanics

Candidates should answer Questions 1 and 2 (10 marks each),  
**and either** Question 3 **or** Question 4 (30 marks).

The content of this sample exam derives from real questions, but the result is  
in many cases test gibberish.

---

**Answer each question in a separate booklet**

Candidates are reminded that devices able to store or display text or images  
may not be used in examinations without prior arrangement.

Approximate marks are indicated in brackets as a guide for candidates.

## SECTION I

- 1 First, *admire* the restful picture of a spiral in Fig. 1, included as a graphic. Fully zenned up? Then let us begin. . . .




Figure 1: A spiral

- (a) Show that, under the action of gravity alone, the scale size of the Universe varies according to

$$\ddot{R} = -\frac{4\pi G\rho_0}{3R^2} \quad (1) \quad [4]$$

and that, consequently,

$$\dot{R}^2 = -\frac{8\pi G\rho_0}{3R} = -K. \quad [3]$$

Express  $K$  in terms of the present values of the Hubble constant  $H_0$  and of the density parameter  $\Omega_0$ . [3]

- (b) In the early Universe, the relation between time and temperature has the form

$$t = \sqrt{\frac{3c^2}{16\pi G g_{\text{eff}} a}} \frac{1}{T^2},$$

where  $a$  is the radiation constant. Discuss the assumptions leading to this equation, but do not carry out the mathematical derivation. Discuss the meaning of the factor  $g_{\text{eff}}$ , and find its value just before and after annihilation of electrons and positrons. [6]

## Q 1 continued

(c) Explain how the present-day neutron/proton ratio was established by particle interactions in the Early Universe. How is the ratio of deuterium to helium relevant to the nature of dark matter? It is *crucially vital* to note that Table 1 is of absolutely no relevance to this question.

Column 1	and row 1
More content	in row 2

Table 1: A remarkably dull table

Finis.

---

Hubble's law:  $v = H_0 D$

---

[4]

[Total: 20]

- 2 (a) The recently-launched *Swift* Gamma Ray Burst telescope is expected to detect about 200 bursts of gamma rays during its 2-year lifespan. Explain why the Poisson distribution,

$$P(n|\lambda) = \exp(-\lambda)\lambda^n/n!$$

is appropriate to describe the probability of detecting  $n$  bursts, and carefully explain the significance of the parameter  $\lambda$ . Table 2 has absolutely nothing to do with this question, and its presence here is proof positive of the existence of aliens who wish to do us typographical harm.

[4]

left	right
------	-------

Table 2: This is a table

Given the above, estimate the probability that *Swift* will detect more than three bursts on any particular calendar day. Blah. Blah. Blaah. Fill the line.

[6]

## Q 2 continued

(b) Explain how Bayesian inference uses the observed number of bursts to infer the true burst rate at the sensitivity limit of *Swift*, and explain the significance of the posterior probability distribution for  $\lambda$ . [5]

Assuming that the posterior,  $p$ , for  $\lambda$  can be approximated as a gaussian, show that, quite generally, the uncertainty in  $\lambda$  inferred from *Swift* will be

$$\sigma \simeq \left( -\frac{\partial^2 \ln p}{\partial \lambda^2} \Big|_{\lambda_0} \right)^{-1/2},$$

where  $\lambda_0$  is the most probable value of  $\lambda$ . [5]

[Total: 20]

3 (a) An earth satellite in a highly eccentric orbit of (constant) perigee distance  $q$  undergoes a tangential velocity impulse  $-\Delta V$  at each perigee passage. By considering the mean rate of change of velocity at perigee, show that the mean rate of change of the semi-major axis  $a$  ( $\gg q$ ) satisfies

$$\frac{1}{a^2} \frac{da}{dt} = \left( \frac{8}{GMq} \right)^{1/2} \frac{\Delta V}{T},$$

where  $M$  is the Earth's mass and  $T$  the orbital period. [3]

---

You may assume  $v^2(r) = GM \left( \frac{2}{r} - \frac{1}{a} \right)$ .

---

Using  $T = 2\pi(a^3/GM)^{1/2}$  show that with  $a_0 = a(0)$ , (where  $a(t)$  is the semimajor axis at time  $t$ )

$$\frac{a(t)}{a_0} = \left[ 1 - \frac{t\Delta V}{2^{1/2}\pi a_0(1-e_0)^{1/2}} \right]^2 \quad [2]$$

and

$$\frac{T(t)}{T_0} = \left[ 1 - \frac{t\Delta V}{2^{1/2}\pi a_0(1-e_0)^{1/2}} \right]^3 \quad [1]$$

and the eccentricity satisfies (with  $e_0 = e(0)$ )

$$e(t) = 1 - \frac{1 - e_0}{\left[ 1 - \frac{t\Delta V}{2^{1/2}\pi a_0(1-e_0)^{1/2}} \right]^2}. \quad [2]$$

Show that, once the orbit is circular, its radius decays exponentially with time on timescale  $m_0/2\dot{m}$  where  $m_0$  is the satellite mass and  $\dot{m}$  the mass of atmosphere 'stopped' by it per second. [2]



## SECTION II

- 4 Show by considering the Newtonian rules of vector and velocity addition that in Newtonian cosmology the cosmological principle demands Hubble's Law  $v_r \propto r$ . [10]

Prove that, in Euclidean geometry, the number  $N(F)$  of objects of identical luminosity  $L$ , and of space density  $n(r)$  at distance  $r$ , observed with radiation flux  $\geq F$  is (neglecting other selection and redshift effects)

$$N(F) = 4\pi \int_0^{(\frac{L}{4\pi F})^{1/2}} n(r)r^2 dr. \quad [5]$$

Use this to show that for  $n = n_1 = \text{constant}$  at  $r < r_1$  and  $n = n_2 = \text{constant}$  at  $r > r_1$ ,

$$N(F) = N_1 \left( \frac{F}{F_1} \right)^{-3/2} \quad \text{for } F > F_1,$$

and

$$N(F) = N_1 \left\{ 1 + \frac{n_2}{n_1} \left[ \left( \frac{F}{F_1} \right)^{-3/2} - 1 \right] \right\} \quad \text{for } F < F_1,$$

where  $F_1 = L/4\pi r_1^2$ ,  $N_1 = N(F_1) = \frac{4}{3}\pi r_1^3 n_1$ . [9]

Reduce these two expressions to the result for a completely uniform density universe with  $n_1 = n_2 = n_0$ . [3]

Sketch how  $n(F)$  would look in universes which are

- flat,
- open,
- and closed. [3]

[Total: 30]

### Cosmology question number 3

- 5 The Friedmann equations are written, in a standard notation,

$$H^2 = \frac{8\pi G\rho}{3} - \frac{kc^2}{R^2} + \frac{\Lambda}{3},$$

$$\frac{d}{dt}(\rho c^2 R^3) = -p \frac{dR^3}{dt},$$

### Q 5 continued

Discuss briefly the meaning of each of  $H$ ,  $\rho$ ,  $k$  and  $\Lambda$ . [4]

Suppose the Universe consists of a single substance with equation of state  $p = w\rho c^2$ , where  $w = \text{constant}$ . Consider the following cases, with  $k = \Lambda = 0$ :

(a) For  $w = 0$ , find the relation between  $R$  and  $\rho$ . Hence show that  $H = \frac{2}{3t}$ . What is the physical interpretation of this case? [8]

(b) In the case  $w = -1$ , show that  $H = \text{constant}$  and  $R = A \exp(Ht)$ , with  $A$  constant. [4]

(c) Explain how the case,  $w = -1$ ,  $k = \Lambda = 0$ ,  $\rho = 0$  is equivalent to an empty, flat, Universe with a non-zero  $\Lambda$ . [2]

(d) Consider a model Universe which contained matter with equation of state with  $w = 0$  for  $0 < t < t_0$ , but which changes to  $W = 0$  for  $t \geq t_0$  without any discontinuity in  $H(t)$ . Regarding this second stage as driven by a non-zero  $\Lambda$  what is the value of  $\Lambda$  if  $t_0 = 10^{24} \mu\text{s}$ ? Define the dimensionless deceleration parameter,  $q$ , and find its value before and after  $t_0$ . **Shout it loud: I'm a geek and I'm proud** [8]

Note: that's

$$t_0 = 10^{24} \mu\text{s} \quad \text{with a letter mu: } \mu.$$

(e) To what extent does this idealized model resemble the currently accepted picture of the development of our Universe? [4]

[Total: 30]

6 In 1908, where was there an airburst 'impact'?

- A. Tunguska
- B. Arizona
- C. Off the Mexican coast
- D. Swindon

- 7 The fossil record suggests that mass extinction events occur once every how many years?
- A. 2.6 Billion Years
  - B. 260 Million Years
  - C. 26 Million Years
  - D. 4 Thousand Years after the dominant lifeform invents fire
- 8 The habitable zone of our Solar system extends over what distances from the Sun?
- A. 0.6–1.5 AU
  - B. 6–15 AU
  - C. 60–150 AU
  - D. 600–1500 AU
  - E. From the little bear's bed all the way through to daddy bear's bed. This is known as the 'Goldilocks zone'.
- 9 If the temperature of the Sun were to increase by 10%, how would the position of the solar habitable zone change?
- A. It would move closer to the Sun.
  - B. It would move further from the Sun.
  - C. It would move to Stornoway.



### SECTION III

- 99 Two variables,  $A$  and  $B$ , have a joint Gaussian probability distribution function (pdf) with a negative correlation coefficient. Sketch the form of this function as a contour plot in the  $AB$  plane, and use it to distinguish between the most probable joint values of  $(A, B)$  and the most probable value of  $A$  given (a different)  $B$ . [5]

Note that this is question 99 on p.9.

- Explain what is meant by *marginalisation* in Bayesian inference and how it can be interpreted in terms the above plot. [5]

Doppler observations of stars with extrasolar planets give us data on  $m \sin i$  of the planet, where  $m$  is the planet's mass and  $i$  the angle between the normal to the planetary orbit and the line of sight to Earth (i.e. the orbital inclination), which can take a value between 0 and  $\pi/2$ .

- Assuming that planets can orbit stars in any plane, show that the probability distribution for  $i$  is  $p(i) = \sin i$ . [5]

A paper reports a value for  $m \sin i$  of  $x$ , subject to a Gaussian error of variance  $\sigma^2$ . Assuming the mass has a uniform prior, show that the posterior probability distribution for the mass of the planet is

$$p(m|x) \propto \int_0^1 \exp \left[ -\frac{(x - m\sqrt{1 - \mu^2})^2}{2\sigma^2} \right] d\mu,$$

- where  $\mu = \cos i$ . [9]

- Determine the corresponding expression for the posterior pdf of  $\mu$ , and explain how both are normalised. [6]

[Total: 30]

- 11 Distinguish between frequentist and Bayesian definitions of probability, and explain carefully how parameter estimation is performed in each regime. [10]

Note that this is question 11 on p.10. It's the one after question 99.

A square ccd with  $M \times M$  pixels takes a dark frame for calibration purposes, registering a small number of electrons in each pixel from thermal noise. The probability of there being  $n_i$  electrons in the  $i$ th pixel follows a Poisson distribution, i.e.

$$P(n_i|\lambda) = \exp(-\lambda)\lambda^{n_i}/n_i!,$$

where  $\lambda$  is the same constant for all pixels. Show that the expectation value of is  $\langle n_i \rangle = \lambda$ . [5]

[You may assume the relation  $\sum_0^\infty \frac{x^n}{n!} = \exp(x)$ .]

Show similarly that

$$\langle n_i(n_i - 1) \rangle = \lambda^2.$$

and hence, or otherwise, that the variance of  $n_i$  is also  $\lambda$ . [5]

The pixels values are summed in columns. Show that these sums,  $S_j$ , will be drawn from a parent probability distribution that is approximately

$$p(S_j|\lambda) = \frac{1}{\sqrt{2\pi M\lambda}} \exp\left[-\frac{(S_j - M\lambda)^2}{2M\lambda}\right],$$

clearly stating any theorems you use. [5]

Given the set of  $M$  values  $\{S_j\}$ , and interpreting the above as a Bayesian likelihood, express the posterior probability for  $\lambda$ , justifying any assumptions you make. [5]

[Total: 30]

## SECTION IV

- 12 Give the equations of motion for  $i = 1, \dots, N$  particles of masses  $m_i$  and positions  $r_i(t)$  under the action of mutual gravity alone in an arbitrary inertial frame. [4]

Use these to derive the following conservation laws of the system:

(a) Constancy of linear momentum – i.e., centre of mass fixed in a suitable inertial frame. [4]

(b) Constancy of angular momentum. [6]

**Q 12 continued**

(c) Constancy of total energy. [8]

How many integrals of motion exist in total? [2]

Derive the moment of inertia of the system and demonstrate its relevance to criteria for escape of particles from the system. [6]

[Total: 30]

**13** For a system of  $N$  objects, each having mass  $m_i$  and position vector  $\mathbf{R}_i$  with respect to a fixed co-ordinate system, use the moment of inertia

$$I = \sum_{i=1}^N m_i R_i^2$$

to deduce the virial theorem in the forms

$$\ddot{I} = 4E_k + 2E_G = 2E_k + 2E$$

where  $E_k$  and  $E_G$  are respectively the total kinetic and gravitational potential energy, and  $E$  is the total energy of the system. [8]

Given the inequality

$$\left( \sum_{i=1}^N a_i^2 \right) \left( \sum_{i=1}^N b_i^2 \right) \geq \left( \sum_{i=1}^N \mathbf{a}_i \cdot \mathbf{b}_i \right)^2 + \left( \sum_{i=1}^N \mathbf{a}_i \times \mathbf{b}_i \right)^2$$

for arbitrary vectors  $\mathbf{a}_i, \mathbf{b}_i, i = 1, \dots, N$ , deduce the following relationship for the  $N$ -body system

$$\frac{1}{4} \dot{J}^2 + J^2 \leq 2IE_k,$$

where  $\mathbf{J}$  is the total angular momentum of the system. [8]

Assuming the system is isolated, use the virial theorem to deduce further the generalised Sundman inequality

$$\frac{\dot{\sigma}}{\rho} \geq 0,$$

in which  $\rho^2 = I$  and  $\sigma = \rho \dot{\rho}^2 + \frac{J^2}{\rho} - 2\rho E$ . [8]

Why does this inequality preclude the possibility of an  $N$ -fold collision for a system with finite angular momentum? [6]

[Total: 30]

**End of Paper**

NOTE: Shout it loud: I'm a geek and I'm proud

NOTE: No correct MCQ answer provided in question 7

NOTE: Too many correct MCQ answers provided in question 8

NOTE: Too few potential answers in MCQ 9