

Garamond-Math, Ver. 2019-08-16

Yuansheng Zhao, Xiangdong Zeng

November 16, 2019

1 Introduction

Garamond-Math is an open type math font matching the *EB Garamond (Octavio Pardo)*¹ and *EB Garamond (Georg Mayr-Duffner)*². Many mathematical symbols are derived from other fonts, others are made from scratch. The metric is generated with a python script.

The font is mostly tested with X_YTeX, though it should also work with LuaTeX.

Issues, bug reports, forks and other contributions are welcome. Please visit GitHub³ for development details.

A minimal example with `unicode-math` package is as following:

```
%Compile with 'xelatex' command
\documentclass{article}
\usepackage[math-style=ISO, bold-style=ISO]{unicode-math}
\setmainfont{EB Garamond}%You should have installed the font
\setmathfont{Garamond-Math.otf}[StylisticSet={7,9}]%Use StylisticSet that you like
\begin{document}
  \[x^3+y^3=z^3.\]
\end{document}
```

The result should be

$$x^3 + y^3 = z^3.$$

2 Alphabets & StylisticSets

Latin and Greek (StylisticSet 4/5 give semi/extra bold for `\symbf`)

ABCDEFGHIJKLMN OPQRSTUVWXYZ

abcdefghijklmnopqrstu vwxyz

ABCDEFGHIJKLMN OPQRSTUVWXYZ

abcdefghijklmnopqrstu vwxyz

ABCDEFGHIJKLMN OPQRSTUVWXYZ

abcdefghijklmnopqrstu vwxyz

ABCDEFGHIJKLMN OPQRSTUVWXYZ

abcdefghijklmnopqrstu vwxyz

ΑΒΓΔΕΖΗΘΘΙΚΛΜΝΞΟΠΡΣΤΥΦΧΨΩ

αβγδεεζήθθικκλμνξοππρρςςττυφφχψω

ΑΒΓΔΕΖΗΘΘΙΚΛΜΝΞΟΠΡΣΤΥΦΧΨΩ

αβγδεεζήθθικκλμνξοππρρςςττυφφχψω

ΑΒΓΔΕΖΗΘΘΙΚΛΜΝΞΟΠΡΣΤΥΦΧΨΩ

¹<https://ctan.org/pkg/ebgaramond/>, and <https://github.com/octaviopardo/EBGaramond12/>

²<https://github.com/georgd/EB-Garamond/>

³<https://github.com/YuanshengZhao/Garamond-Math/>

αβγδεεζηθθικκλμνξοππρρσςτυφφχψω
ΑΒΓΔΕΖΗΘΘΙΚΑΜΝΞΟΠΡΣΤΥΦΧΨΩ
αβγδεεζηθθικκλμνξοππρρσςτυφφχψω
ΑΒCDEFGHIJKLMNOPQRSTUVWXYZ
abcdefghijklmnopqrstuvwxy
ΑΒCDEFGHIJKLMNOPQRSTUVWXYZ
abcdefghijklmnopqrstuvwxy

Sans and Typewriter: From Libertinus Math⁴

ΑΒCDEFGHIJKLMNOPQRSTUVWXYZ
abcdefghijklmnopqrstuvwxy
ΑΒCDEFGHIJKLMNOPQRSTUVWXYZ
abcdefghijklmnopqrstuvwxy
ΑΒCDEFGHIJKLMNOPQRSTUVWXYZ
abcdefghijklmnopqrstuvwxy
ΑΒCDEFGHIJKLMNOPQRSTUVWXYZ
abcdefghijklmnopqrstuvwxy
ΑΒCDEFGHIJKLMNOPQRSTUVWXYZ
abcdefghijklmnopqrstuvwxy

Blackboard (StylisticSet 1 → rounded XITS Math⁵)

ΑΒCDEFGHIJKLMNOPQRSTUVWXYZ
abcdefghijklmnopqrstuvwxy
ΑΒCDEFGHIJKLMNOPQRSTUVWXYZ
abcdefghijklmnopqrstuvwxy

Script: Rounded XITS Math [StylisticSet 3 → scaled CM; 8 → Garamond-compatible ones (experimental)]

ΑΒCDEFGH IJKL MNOP QRSTUVWXYE
abcdefghijklmnopqrstuvwxy
ΑΒCDEFGH IJKL MNOP QRSTUVWXYE
abcdefghijklmnopqrstuvwxy
ΑΒCDEFGHIJKLMNOPQRSTUVWXYZ
ΑΒCDEFGHIJKLMNOPQRSTUVWXYZ
ΑΒCDEFGH IJKL MNOP QRSTUVWXYE
abcdefghijklmnopqrstuvwxy

Fraktur: From Noto Sans Math⁶

ΑΒCDEFGH IJKL MNOP QRSTUVWXYE
abcdefghijklmnopqrstuvwxy
ΑΒCDEFGH IJKL MNOP QRSTUVWXYE
abcdefghijklmnopqrstuvwxy

⁴<https://github.com/khaledhosny/libertinus/>
⁵<https://github.com/khaledhosny/xits/>
⁶<https://github.com/googlefonts/noto-fonts/>

Digits: Same width between weight and serif/sans

3.141592653589793238462643383279502884197169399375105820974944592307816406286
3.141592653589793238462643383279502884197169399375105820974944592307816406286
3.141592653589793238462643383279502884197169399375105820974944592307816406286

\partial: (StylisticSet 2 → curved ones)

$$\partial_\mu(\partial^\mu\phi) - \epsilon^{\lambda\mu\nu}\partial_\mu(A_\lambda\partial_\nu f)$$
$$\partial_\mu(\partial^\mu\phi) - \epsilon^{\lambda\mu\nu}\partial_\mu(A_\lambda\partial_\nu f)$$

\hbar: (StylisticSet 6 → horizontal bars)

\hbar \hbar

Italic \hbar : (StylisticSet 10 → out-bending ones)

$$\hbar = \frac{\hbar}{2\pi} \quad \hbar = \frac{\hbar}{2\pi}$$

\tilde: (StylisticSet 9 → “normal” ones)

\tilde{F} \tilde{F}

\int: (StylisticSet 7 → a variant with inversion symmetry)

$$\oint_{\partial\Sigma} \vec{E} \cdot d\vec{l} = -\frac{1}{c} \frac{d}{dt} \iint_{\Sigma} \vec{B} \cdot d\vec{S}$$
$$\oint_{\partial\Sigma} \vec{E} \cdot d\vec{l} = -\frac{1}{c} \frac{d}{dt} \iint_{\Sigma} \vec{B} \cdot d\vec{S}$$

Binany Operators: (StylisticSet 11 → larger ones)


$$s = A + b \times 1 \div x^3$$
$$s = A + b \times 1 \div x^3$$

Extensible Arrow Hack

The font contains the math table for constructing extensible arrow. However `unicode-math` does not provide an interface to that. In `LuaTeX` one can use `\Uhexensible`⁷. A more general solution is to add the following code in preamble.

```
\usepackage{extarrow} %or mathtools
\makeatletter
\renewcommand{\relbar}{\symbol{"E010}\mkern-.2mu\symbol{"E010}\mkern1.8mu}
\renewcommand{\Relbar}{\symbol{"E011}\mkern-.2mu\symbol{"E011}\mkern1.8mu}
\makeatother
```

Then `\xleftarrow` and other commands will work:



3 Known Issue

- Various spacing problems. Though math fonts technically should not be kerned, some pairs look very ugly (Ex. \mathcal{VA}); sometimes sub/superscript may also have same problem. However, do note that due to the mechanism in math mode, making all spacing look perfect is almost impossible (as far as I can do, and low x-height and large italic angle only make things even worse), in many cases, adjusting manually (i.e. using `\`, or `\!`) is required.
- Fake optical size. EB Garamond does not contain a complete set of glyphs (normal + bold + optical size of both weights). The “optical size `ss ty`” is made by interpolating different weights at the present (without this, the double script is too thin to be readable).

⁷<https://tex.stackexchange.com/questions/423893/>

4 Equation Samples

$$1 + 2 - 3 \times 4 \div 5 \pm 6 \mp 7 + 8 = -a \oplus b \otimes c - \{z\}$$

$$\forall \epsilon, \exists \delta : x \in A \cup B \subset S \cap T \not\subseteq U$$

$$R_{\nu\kappa\lambda}^{\mu} = \partial_{\kappa} \Gamma_{\lambda\nu}^{\mu} - \partial_{\lambda} \Gamma_{\kappa\nu}^{\mu} + \Gamma_{\kappa\sigma}^{\mu} \Gamma_{\lambda\nu}^{\sigma} - \Gamma_{\lambda\sigma}^{\mu} \Gamma_{\kappa\nu}^{\sigma}$$

$$T_{\alpha_1 \dots \alpha_k}^{\beta_1 \dots \beta_l} = T_{i_1 \dots i_k}^{j_1 \dots j_l} \frac{\partial x^{i_1}}{\partial x'^{\alpha_1}} \dots \frac{\partial x^{i_k}}{\partial x'^{\alpha_k}} \frac{\partial x'^{\beta_1}}{\partial x^{j_1}} \dots \frac{\partial x'^{\beta_l}}{\partial x^{j_l}}$$

$$\int_{\sqrt{\frac{1-mu+md}{2mn/k}}}^{X_p} \overbrace{1+2+3+4+5+6+7+8}$$

$$x \leftarrow y \leftrightarrow w \Rightarrow b \Leftrightarrow c \uparrow y \updownarrow w \Downarrow b \Downarrow c \searrow p \not\ll px \leftarrow x \uparrow X \leftarrow Y \mapsto Z \uparrow f \rightleftharpoons f \updownarrow fb \Rrightarrow b \Leftrightarrow p$$

$$\int_0^1 \frac{\ln(x+1)}{x} dx = \int_0^1 \sum_{i=1}^{\infty} \frac{(-x)^{i-1}}{i} dx = \sum_{i=1}^{\infty} \int_0^1 \frac{(-x)^{i-1}}{i} dx = \sum_{i=1}^{\infty} \frac{(-1)^{i+1}}{i^2} = \frac{\pi^2}{12}$$

$$\int_0^{\infty} \int_0^{\infty} \sum_{i=1}^{\infty} \prod_{j=i}^{\infty} \prod_{k=i}^{\infty} \text{ff} \text{ff} \text{ff} \text{ff}$$

$$\left(\left(\left((x) \right) \right) \right) \left[\left[\left[[x] \right] \right] \right] \left\{ \left\{ \left\{ \{x\} \right\} \right\} \right\} \left| \left| \left| |x| \right| \right| \right| \left| \left| \left| \|x\| \right| \right| \right| \left| \left| \left| \langle x \rangle \right| \right| \right|$$

$$\left(\left(\left((x) \right) \right) \right) \left[\left[\left[[x] \right] \right] \right] \left[\left[\left[[x] \right] \right] \right]$$

$$\langle x | + |x\rangle + \langle \alpha | \beta \rangle + |\alpha\rangle \langle \beta| + \left\langle \frac{1}{2} \right| + \left| \frac{1}{2} \right\rangle + \left\langle \frac{1}{2} \right| \frac{1}{2} \rangle + \left| \frac{1}{2} \right\rangle \left\langle \frac{1}{2} \right| + \left\langle \frac{a^2}{b^2} \right| + \left| \frac{e^{x^2}}{e^{y^2}} \right\rangle$$

$$\bullet \bullet \bullet \bullet \bullet \bullet \bullet \bullet \bullet \bullet \bullet \bullet + ABC^{\circledast \circledast \circledast \circledast \circledast \circledast \circledast \circledast \circledast \circledast}$$

$$\begin{pmatrix} u_0 \\ u_1 \\ \vdots \\ u_{N-1} \end{pmatrix} = \sum_{k>0} \left[\begin{pmatrix} 1 \\ \cos ka \\ \vdots \\ \cos k(N-1)a \end{pmatrix} \frac{C_{k+} \cos(\omega_k t + \varphi_{k+})}{\frac{2}{\sqrt{N}} q_{k+}} + \begin{pmatrix} 0 \\ \sin ka \\ \vdots \\ \sin k(N-1)a \end{pmatrix} \frac{C_{k-} \cos(\omega_k t + \varphi_{k-})}{\frac{2}{\sqrt{N}} q_{k-}} \right]$$

$$\begin{aligned} \mathcal{F}^{-1}(|j\rangle) &= \frac{1}{\sqrt{2^n}} \sum_{k=0}^{2^n-1} \exp\left(-2\pi i \frac{jk}{2^n}\right) |k\rangle \\ &= \frac{1}{\sqrt{2^n}} \sum_{k_{n-1}=0}^1 \dots \sum_{k_0=0}^1 \exp\left(-2\pi i j \sum_{l=0}^{n-1} \frac{2^l k_l}{2^n}\right) |k_{n-1} \dots k_0\rangle \\ &= \frac{1}{\sqrt{2^n}} \sum_{k_{n-1}=0}^1 \dots \sum_{k_0=0}^1 \bigotimes_{l=1}^n \left[\exp\left(-2\pi i j \frac{k_{n-l}}{2^l}\right) |k_{n-l}\rangle \right] \\ &= \frac{1}{\sqrt{2^n}} \bigotimes_{l=1}^n \left[\sum_{k_{n-l}=0}^1 \exp\left(-2\pi i j \frac{k_{n-l}}{2^l}\right) |k_{n-l}\rangle \right] \\ &= \frac{1}{\sqrt{2^n}} \bigotimes_{l=1}^n \left[|0\rangle_{n-l} + e^{-2\pi i j / 2^l} |1\rangle_{n-l} \right] \\ &= \frac{1}{\sqrt{2^n}} \bigotimes_{l=1}^n \left[|0\rangle_{n-l} + e^{-2\pi i (0 \cdot j_{l-1} \dots j_0)} |1\rangle_{n-l} \right]. \end{aligned}$$

$$\begin{aligned}
S &= \frac{m}{2} \int_0^{t_f} \left[\left(-\omega x_i \sin \omega t + \omega \frac{x_f - x_i \cos \omega t_f}{\sin \omega t_f} \cos \omega t \right)^2 + \sum_{n=1}^{\infty} \left(\frac{a_n n \pi}{t_f} \right)^2 \cos^2 \frac{n \pi t}{t_f} \right] dt \\
&\quad - \frac{m \omega^2}{2} \int_0^{t_f} \left[\left(x_i \cos \omega t + \frac{x_f - x_i \cos \omega t_f}{\sin \omega t_f} \sin \omega t \right)^2 + \sum_{n=1}^{\infty} a_n^2 \sin^2 \frac{n \pi t}{t_f} \right] dt \\
&= \sum_{n=1}^{\infty} \int_0^{t_f} \left[\frac{m}{2} \left(\frac{a_n n \pi}{t_f} \right)^2 \cos^2 \frac{n \pi t}{t_f} - \frac{m \omega^2}{2} a_n^2 \sin^2 \frac{n \pi t}{t_f} \right] dt \\
&\quad + \frac{m \omega^2}{2} \int_0^{t_f} \left[x_i^2 - \left(\frac{x_f - x_i \cos \omega t_f}{\sin \omega t_f} \right)^2 \right] (\sin^2 \omega t - \cos^2 \omega t) dt \\
&\quad - \frac{m \omega^2}{2} \int_0^{t_f} 4 x_i \left(\frac{x_f - x_i \cos \omega t_f}{\sin \omega t_f} \right) (\sin \omega t \cos \omega t) dt.
\end{aligned}$$

$$\begin{aligned}
U(x_f, t_f; x_i, t_i) &= \sqrt{\frac{m \omega}{2 \pi i \hbar \sin [\omega (t_f - t_i)]}} \\
&\quad \times \exp \left\{ \frac{i m \omega}{2 \hbar \sin [\omega (t_f - t_i)]} \left[(x_i^2 + x_f^2) \cos [\omega (t_f - t_i)] - 2 x_i x_f \right] \right\}.
\end{aligned}$$